

***Bahman Guyuron, MD, FACS***  
***Plastics and Reconstructive Surgery***

**440-461-7999**

**HOME INSTRUCTIONS FOR PTOSIS CORRECTION AND LOWER  
BLEPHAROPLASTY**

- You will have an incision in the crease of the Upper eyelid; Use the Tobradex eye drop three times a day as directed. Use the eye rewetting drops as often as needed.
- Swelling and bruising are expected and will usually peak at 48 hours and will gradually decrease over the next 7-10 days.
- **If the swelling is causing a pressure-type pain to the eyeball, please contact our office immediately.**
- **If at anytime you feel a sharp pain or sensation of a foreign body in the eye, please contact the office immediately.**
- You may have food as tolerated, but avoid highly salty foods.
- Sleep on your back with your head elevated at approximately a 45 degree angle.
- You may wash your face with a washcloth except for the eye area.
- Avoid bending over for long periods of time.
- Sunglasses are recommended when outdoors
- Avoid strenuous activity for three weeks post operatively. Slow walking is encouraged immediately after surgery. Rotate your ankles frequently while lying in bed.
- Do not wear clothing that pulls over the head.
- Do not take any aspirin, Advil or Aleve for a period of one week following your surgery.
- If you need pain medication, do not hesitate to take this as prescribed. However, **please eat food immediately after taking the medication to minimize the chance for nausea or vomiting.**
- **All patients are seen the day after surgery.**
- **Someone must be with you for 24 hours after surgery.**
- You may not be able to drive yourself for 5 to 7 days after surgery if you are taking pain medications, please arrange for your transportation for the post operative appointments.

**REMINDERS:**

- Refer to the patient discharge Instructions upon discharge from the surgery center.
- Please take all of your medications on schedule
- Report any unusual heavy bleeding or discomfort unless you have been advised otherwise. Report shortness of breath, temperature above 100 degrees and redness. Pain or discomfort in your legs should be reported immediately.

**EMERGENCY NUMBERS:**

For emergencies after normal office hours, please call the answering service at **440-461-7999**. Leave your name, phone number and the reason for your call. If there is not a response within 10 minutes, please call the number again.